

Tutorial Problems #5

MAT 267 – Advanced Ordinary Differential Equations – Fall 2014

Christopher J. Adkins

SOLUTIONS

Reduction of Order Via Differential Operators Let $D = \frac{d}{dx}$ be our differential operator. Then any n -th order linear non-homogeneous equation may be written as

$$L(D)[y(x)] = f(x) \quad \text{where} \quad L(D) = D^n + a_{n-1}D^{n-1} + \dots + a_0 \quad \left(D^n = \frac{d^n}{dx^n} \right)$$

with $a_i \in \mathbb{R}$. Factor L into a product of its roots (which may be complex and we'll deal with later), i.e.

$$L(D) = (D - \lambda_1) \dots (D - \lambda_n)$$

Notice this factorization is not possible if a_i are functions since the differential operator isn't commutative ($D_1D_2 = D_2D_1$). Thus, if we let $y_n = (D - \lambda_n)y$ and $y_i = (D - \lambda_i)y_{i+1}$ we effectively reduce the n -th order equations into n first order equations (which we know how to handle)

pg. 267 - # 28 Solve (using reduction of order)

$$y'' + y' = x^2 + 2x$$

Solution We see that if $L = D^2 + D$, then

$$L(D)[y(x)] = x^2 + 2x$$

is the ODE we're looking to solve. Notice we may use the above method to deduce

$$L(D) = D(D + 1) \implies u' = x^2 + 2x \quad \text{where} \quad (D + 1)y = u$$

The above ODE in u is separable, thus

$$u(x) = \int x^2 + 2x dx = \frac{x^3}{3} + x^2 + C_1 \quad C_1 \in \mathbb{R}$$

We now know

$$y' + y = \frac{x^3}{3} + x^2 + C_1$$

which is a first order linear ODE, we know this may be solved using an integrating factor. We know

$$y(x) = \frac{1}{\mu(x)} \int \mu(x)g(x)dx \quad \text{where} \quad \mu(x) = \exp\left(\int dx\right) = e^x$$

Thus the general solution to the ODE is

$$y(x) = e^{-x} \int e^x \left(\frac{x^3}{3} + x^2 + C_1 \right) dx = \frac{x^3}{3} + C_2 e^{-x} + C_1 \quad C_2 \in \mathbb{R}$$

□

The Inverse of a Differential Operator Let's talk about D^{-1} now. Formally we need an operator with the property if $Dx = y$, then $x = D^{-1}y$. Intuitively, you should think the integral operator is a natural left inverse for D since

$$\frac{d}{dx} \int f(x) dx = f(x)$$

by the fundamental theorem of calculus. Now what about factors of $(D - \lambda)$ we had...using a formal series expansion (notably a geometric series), we may algebraically write

$$(D - \lambda)^{-1} = -\frac{1}{\lambda(1 - D/\lambda)} = -\frac{1}{\lambda} \left[1 + \frac{D}{\lambda} + \frac{D^2}{\lambda^2} + \frac{D^3}{\lambda^3} \dots \right]$$

Convergence of this series is a slight issue at the moment...but for any solution that terminates after a finite number of derivatives we know convergence is guaranteed. Let's revisit the example we just saw.

pg. 267 - # 28 Solve (using Inverse Operators)

$$y'' + y' = x^2 + 2x$$

Solution As we saw before we have

$$D(D + 1)y = x^2 + 2x \implies y_p(x) = \frac{1}{D(D + 1)}(x^2 + 2x)$$

Notice we'll only be able to pick up the particular solution to the ODE with this method (not the general) since L is not injective in general (i.e. $L[y_{hom}] = 0$). Expanding the inverse into formal series shows

$$y_p(x) = \frac{1}{D} (1 + D + D^2) (x^2 + 2x) = \left[\frac{1}{D} - 1 + D \right] (x^2 + 2x)$$

Thus

$$y_p(x) = \int (x^2 + 2x) dx - (x^2 + 2x) + \frac{d}{dx} (x^2 + 2x) = \frac{x^3}{3} + 2$$

You may recover the general solution using your knowledge of homogeneous equation, but seeing the eigenvalues of $\lambda = 0$ and $\lambda = -1$, thus 1 and e^{-x} solve the homogeneous problem. □

A Special Case, $(D - \lambda)^{-1}$ Applied To e^{ax} Notice in the case of exponential, we may factor out e^{ax} in the formal expansion since

$$\frac{d^n}{dx^n} e^{ax} = a^n e^{ax}$$

Thus

$$\frac{1}{D - \lambda} e^{ax} = -\frac{e^{ax}}{\lambda} \left[1 + \frac{a}{\lambda} + \frac{a^2}{\lambda^2} + \dots \right] = \frac{e^{ax}}{a - \lambda}$$

where we side-stepped the notion of convergence once again, but clearly this is an inverse since

$$(D - \lambda) \left(\frac{1}{D - \lambda} e^{ax} \right) = (D - \lambda) \frac{e^{ax}}{a - \lambda} = e^{ax}$$

Since this will work with any $a \in \mathbb{C}$ and the inverse raised to integer powers, we've therefore found a way to handle exponentials. The only issue that may occur is if $a = \lambda$ since the expansion isn't defined (in other words, a is an eigenvalue). This can easily be fixed using the exponential shift theorem,

$$L(D)[e^{ax}y] = e^{ax}L(D + a)[y]$$

when $L(x)$ is a polynomial (the proof goes by induction, and also applies to the inverse). Thus if λ is a root of L , i.e. $L(D) = (D - \lambda)^k g(D)$, we see that

$$\frac{1}{(D - \lambda)^k g(D)} e^{\lambda x} = e^{\lambda x} \frac{1}{D^k g(D + \lambda)} 1 = e^{\lambda x} \frac{x^k}{k! g(\lambda)}$$

pg. 282 - # 32 Solve

$$y''' + y' = \cos x$$

Solution Well, in terms of D we have that

$$D(D - i)(D + i)y = \cos x$$

Now since we've just dealt with exponentials so far, note $e^{ix} = \cos x + i \sin x$, so lets solve

$$D(D - i)(D + i)y = e^{ix}$$

and take the real part. Letting $g(D) = D(D + i)$ like the above, we see

$$y_p(x) = \frac{1}{(D - i)g(D)} e^{ix} = e^{ix} \frac{x}{g(i)} = -e^{ix} \frac{x}{2} = -\frac{x \cos x}{2} - i \frac{x \sin x}{2}$$

Since we just want the real part of the solution, we see the particular solution to the ODE is

$$y_p(x) = -\frac{x \cos x}{2}$$

Noting that the eigenvalues of the equation are $\lambda = 0, \pm i$, we have that

$$y(x) = C_1 + C_2 \cos x + C_3 \sin x - \frac{x \cos x}{2}$$

is the general solution. □

Partial Fraction Decomposition with Differential Operators As you've probably seen before with polynomials, you may decompose

$$\frac{1}{(D + \lambda_1)(D + \lambda_2)} = \frac{c_1}{D + \lambda_1} + \frac{c_2}{D + \lambda_2}$$

Let's see how this would apply to the previous example.

pg. 282 - # 32 Solve (using partial fractions)

$$y''' + y' = \cos x$$

Solution Using what we saw before, let's try to decompose into pieces:

$$\frac{1}{D(D-i)(D+i)} = \frac{c_1}{D} + \frac{c_2}{D-i} + \frac{c_3}{D+i}$$

This implies we need

$$(D-i)(D+i)c_1 + D(D+i)c_2 + D(D-i)c_3 = 1 \implies \begin{cases} c_1 + c_2 + c_3 = 0 \\ c_2 - c_3 = 0 \\ c_1 = 1 \end{cases} \implies c_2 = -\frac{1}{2}, c_3 = -\frac{1}{2}$$

Thus

$$\frac{1}{D(D-i)(D+i)} = \frac{1}{D} - \frac{1}{2(D-i)} - \frac{1}{2(D+i)}$$

Now if we apply this to e^{ix} , we see

$$\frac{1}{D(D-i)(D+i)} e^{ix} = \frac{e^{ix}}{i} - \frac{x e^{ix}}{2} - \frac{e^{ix}}{4i} + C = \frac{3}{4i} e^{ix} - \frac{x e^{ix}}{2} + C$$

If we take the real part of this solution we see the following particular solution

$$y_p(x) = \frac{3}{4} \sin x - \frac{x \cos x}{2} + C$$

□

Quiz Find the partial solution using any inverse operator method for

$$y'' + 3y' + 2y = 2(e^{-2x} + x^2)$$

Solution We see that

$$L(D) = (D+2)(D+1)$$

Thus we want to solve

$$y_p(x) = \frac{1}{(D+2)(D+1)} (2e^{-2x} + 2x^2)$$

For the exponential, we may use what we've previous talked about to find that we have $L = (D+2)g(D)$, hence

$$y_{p_e}(x) = \frac{2xe^{-2x}}{g(-2)} = -2xe^{-2x}$$

For the polynomial, we have that

$$y_{p_p}(x) = \frac{1}{(D+2)(D+1)} 2x^2 = \frac{1}{2} \left(1 - \frac{D}{2} + \frac{D^2}{4} \right) (1 - D + D^2) 2x^2 = \frac{1}{2} \left(1 - \frac{3D}{2} + \frac{7D^2}{4} \right) 2x^2$$

Thus we see

$$y_{p_p}(x) = x^2 - 3x + \frac{3}{2}$$

So

$$y_p(x) = -2xe^{-2x} + x^2 - 3x + \frac{7}{2}$$

is a particular solution.

□